

About EAS inverse approach

S. V. Ter-Antonyan*

Yerevan Physics Institute, Alikhanyan Brothers 2, Yerevan 375036, Armenia

Abstract

It is shown that using the observed extensive air shower (EAS) electron and truncated muon size spectra at sea level one can solve the EAS inverse problem - reconstruction of primary energy spectra and elemental composition, for not more than 2 kinds of primary nuclei.

*e-mail:samvel@jerewan1.yerphi.am

1 Introduction

High accuracy of modern EAS experiments in the primary energy region of 10^{15} eV [1-4] increased a number of publications on the solutions of the EAS inverse problem, which is reconstruction of primary nuclei energy spectra based on the detected EAS parameters at observation level [5-9]. However, discrepancies in results continue to grow. In most of the cases, it is a result of the hidden experimental systematic errors due to uncertainties in the response functions of detectors. The uncertainty of A-A interaction model at these energies also contributes to the discrepancies in primary nuclei energy spectra results. However, there are certain publications where the EAS inverse problem is solved based on erroneous presuppositions, which of course only increase the existing discrepancies of data.

2 EAS inverse problem

In general, the relation between energy spectra ($\partial\mathfrak{Z}_A/\partial E$) of primary nuclei ($A \equiv 1, 4, \dots 56$) and measured EAS electron and muon size spectra at observation level ($\Delta I/\Delta N_{e,\mu}^*$) at a given zenith angle θ is determined by an integral equation

$$\frac{\Delta I(\theta)}{\Delta N_{e,\mu}^*} = \sum_A \int \frac{\partial\mathfrak{Z}_A}{\partial E} \frac{\partial K(E, A, \theta)}{\partial N_{e,\mu}^*} dE \quad (1)$$

The kernel function of equation (1) is determined as

$$\frac{\partial K}{\partial N_{e,\mu}^*} \equiv \int \frac{\partial W(E, A, \theta)}{\partial N_{e,\mu}} \frac{dG}{dN_{e,\mu}^*} dN_{e,\mu} \quad (2)$$

where the EAS size spectra $\partial W/\partial N_{e,\mu}$ depend on observation level, A, E, θ parameters of primary nucleus and $A - A_{Air}$ interaction model, $dG(N, N^*)/dN^*$ is the error function of measurements. Equation (1) is a typical ill-posed problem and has an infinite set of solutions for unknown primary energy spectra. However, the integral equation (1) turns to a Fredholm equation if a kind of a primary nucleus is defined directly in the experiment (as it successfully does in the balloon and satellite measurements [10, 11] where energy spectra for different nuclei are obtained up to 10^{15} eV) or the detected EAS parameters (or combination of parameters) slightly depend on primary nuclei [5]. In the last case one evaluate only the all-particle energy spectra $\sum_A d\mathfrak{Z}_A/dE$ from equation (1).

As a result, the meaning of data presented in the [9] and further publications [12-14] where authors claim to have got the solution of equation (1) for 4 types of primary nuclei, is not clear.

Let's prove that using the measured EAS electron and truncated muon size spectra at 3 zenith angular intervals, as its done in [9], the Eq. (1) can have a single solution only for primary flux which consists of not more than 2 kinds of nuclei.

Let the EAS size spectra at observation level KASCADE ($t = 1020$ g/cm² [9]) be described by log-Gaussian distributions with mean $\langle N_{e,\mu}(E, A, \theta) \rangle$ and variance $\sigma_{e,\mu}^2(E, A, \theta)$. It is known that this assumption is well performed at atmosphere depth $t > 700$ g/cm², primary energies $E > 10^5$ GeV and zenith angles $\theta < 35^\circ$.

Let also the measurements and further evaluations of EAS electron and truncated muon sizes be carried out without errors ($dG/dN^* \equiv \delta(N - N^*)$) and integral Eq. (1) include only statistical uncertainties.

Then a set of Eq. (1) transforms into the following:

$$\frac{\Delta I(\theta)}{\Delta N_{e,\mu}} = \sum_A \int \frac{\partial \Im_A}{\partial E} \frac{\partial W(E, A, \theta)}{\partial N_{e,\mu}} dE \quad (3)$$

Let's determine the parameters of distribution functions $\partial W/\partial N_{e,\mu}$ by the following known empirical expressions:

$$\langle N \rangle \simeq a \left(\frac{E}{1 \text{ GeV}} \right)^b A^c \cos^d \theta \quad (4)$$

$$\sigma \simeq \alpha A^\delta \left(\frac{\ln(E/1 \text{ GeV})}{\ln 10^6} \right)^\varepsilon \cos^\rho \theta \quad (5)$$

where the values of corresponding approximation parameters ($a, \dots, d, \alpha, \dots, \rho$) are presented in Tables 1,2 and obtained by CORSIKA6016(NKG) EAS simulation code [15] at QGSJET interaction model [16].

	a	b	c	d
e	0.0041 ± 0.0004	1.22 ± 0.005	-0.22 ± 0.01	6.8 ± 0.2
μ	0.0079 ± 0.0003	0.933 ± 0.001	0.07 ± 0.002	1.87 ± 0.05

Table 1: Approximation parameters of average EAS electron size (N_e) and EAS truncated muon size (N_μ) in empirical formula (4).

	α	δ	ε	ρ
e	0.79 ± 0.01	0.25 ± 0.003	1.82 ± 0.05	2.1 ± 0.1
μ	0.283 ± 0.003	0.277 ± 0.003	1.12 ± 0.05	0.26 ± 0.1

Table 2: Approximation parameters of $\sigma_{e,\mu}$ in empirical formula (4).

The accuracy of approximation (4) for average EAS electron size $\langle N_e(E, A, \theta) \rangle$ is less than 10% at $3 \cdot 10^5 < E < 3 \cdot 10^8$ GeV, $A \equiv 1, \dots, 56$, $\theta < 32^\circ$ and observation level 1020 g/cm². The corresponding accuracies of $\sigma_{e,\mu}$ and average EAS truncated muon size N_μ are less than 1-2%. Changing the variables of kernel functions of Eq. (3) from N_e and N_μ to x_e and x_μ respectively according to

$$x_{e,\mu} \equiv A^\delta \left(\frac{N_{e,\mu}}{\langle N_{e,\mu} \rangle} - 1 \right) + 1 \quad (6)$$

we obtain the following set of Fredholm integral equations

$$\frac{\Delta I(\theta)}{\Delta N_{e,\mu}} = \int f_{e,\mu}(E, A) F_{e,\mu}(E, \theta) dE \quad (7)$$

where

$$f_{e,\mu}(E, A) = \sum_A \frac{\partial \mathfrak{S}_A}{\partial E} A^{\delta_{e,\mu} - c_{e,\mu}} \quad (8)$$

and kernel functions

$$F_{e,\mu}(E, \theta) = \frac{\partial W(E, \theta)}{\partial x_{e,\mu}} \frac{1}{a_{e,\mu} E^{b_{e,\mu}} \cos^{d_{e,\mu}} \theta} \quad (9)$$

independent of kind of primary nuclei.

Examples of distribution functions $\partial W/\partial x$ for EAS electron and truncated muon size spectra are presented Fig. 1,2 respectively. These data were obtained by CORSIKA code for primary proton




Figure 1: EAS electron size spectra for primary proton (empty symbols) and iron (filled symbols) nuclei at energy $E = 3.2$ PeV, zenith angle $\theta < 18^\circ$ and observation level 1020 g/cm^2 . The lines are the corresponding log-Gaussian distributions.

($A \equiv 1$, empty symbols) and iron ($A \equiv 56$, filled symbols) nuclei at energy $E = 3.2 \cdot 10^6 \text{ GeV}$ and zenith angle $\theta < 18^\circ$. The lines in Fig. 1,2 correspond to log-Gaussian distributions which were counted based on known values of mean $\langle x \rangle = 1$ and variances $\sigma_{e,\mu}^2(E, \theta)$.

It is seen, that the distribution functions slightly depend on a kind of a primary nucleus, especially the right-hand sides of distributions. It is an important fact, because the contribution of the left-hand sides of distributions is negligible small at *a priori* steep primary energy spectra ($\partial \mathfrak{S}_A/\partial E \sim E^{-3}$).

Let the functions $f_e^{(j,k)}(E_i)$ and $f_\mu^{(j,k)}(E_i)$ be the solutions of the set of Fredholm equations (7) at $i = 1, \dots, m$ values of primary energies E_i , $j = 1, \dots, n$ kinds of primary nuclei and $k=1,2,3$ zenith angles. Then the values of unknown energy spectra ($\partial \mathfrak{S}_{A_j}/\partial E_i$) at a given set of E_i and different primary nuclei ($A_j \equiv A_1, \dots, A_n$) are determined by the set of linear equations

$$\langle f_e^{(j)}(E_i) \rangle_\theta = \sum_{A=A_1}^{A=A_n} A^{\delta_e - c_e} \frac{\partial \mathfrak{S}_A}{\partial E} \Big|_{E=E_i} : i = 1, \dots, m \quad (10)$$

$$\langle f_{\mu}^{(j)}(E_i) \rangle_{\theta} = \sum_{A=A_1}^{A=A_n} A^{\delta_{\mu}-c_{\mu}} \frac{\partial \mathfrak{S}_A}{\partial E} \Big|_{E=E_i} : i = 1, \dots, m \quad (11)$$

where $\langle f_{e,\mu}^{(j)}(E_i) \rangle_{\theta} = \frac{1}{3} \sum_k f_{e,\mu}^{(j,k)}(E_i)$

Evidently, the single solutions of linear equations (9,10) occur only at $mn \leq 2m$, where mn is a number of unknown $(\partial \mathfrak{S}_{A_j} / \partial E_i)$ and $2m$ is a number of equations (10,11).

Therefore the condition of existence of solutions is $n \leq 2$ and the energy spectra [9] obtained




Figure 2: The same as Fig.1 for EAS truncated muon size.

based on KASCADE EAS electron and truncated muon size spectra for 4 ($A \equiv 1, 4, 16, 56$) primary nuclei no physical meaning. At the same time the all-particle spectrum obtained in [9] have to be approximately right due to the power index in Eq. (11) $\delta_{\mu} - c_{\mu} \ll 1$.

It should be noted that unreliability of solutions [9] is also shown in [17] based on correlation analysis between the observable electron and truncated muon size spectra and unknown energy spectra of primary nuclei.

3 Conclusion

Based on equation (1), the investigation of the EAS inverse problem, which is the reconstruction of energy spectra of primary nuclei by the observable EAS electron and muon size spectra at observation level makes sense only using *a priori* given functions for unknown primary energy spectra with given unknown spectral parameters (so called parameterization of equation (1)) as it is done in [8, 18, 19, 20].

References

- [1] H.O. Klages et al., 9th ISVHECR, Karlsruhe (1996) Nucl. Phys. B (Proc. Suppl.).
- [2] M. Aglietta et al., Astropart. Phys. (10) 1 (1999) 1.
- [3] S.V. Ter-Antonyan et al., Proc. 24th ICRC Rome, **1** (1995) 365.
- [4] M.A.K. Glasmacher et al., Astropart. Phys. 10 (1999) 291.
- [5] A.P.Garyaka et al., J.Phys. G: Nucl. Phys., 28 (2002) 2317.
- [6] A. Chilingarian, S. Ter-Antonyan et al., Nucl. Phys. B, 52B, (1997) 240.
- [7] J.R. Horandel, Proc. 27th ICRC Hamburg **1** (2001) 71.
- [8] R.Glasstetter et al., Proc. 26th ICRC Salt Lake City (1999) **1** 222.
- [9] H. Ulrich, et al., Proc. 27th ICRC, Hamburg, **1** (2001) 97.
- [10] K. Asakimory et al., Ap.J. 502 (1998) 278.
- [11] A.V. Apananeko et al., Proc. 26th ICRC, Salt Lake City, **3**, (1999) 163.
- [12] A.Haungs, 18th ECRS, Moscow (2002) Ep1a.2,
<http://dbserv.npi.msu.su/~18sym/conference/transparencies.phtml>.
- [13] J.R.Horandel et al. (KASCADE collaboration), Aquila, Italy, (2001),
http://www-ik.fzk.de/KASCADE/KASCADE_publications.html
- [14] K.-H. Kampert et al. (KASCADE collaboration), Highlight Paper, Proc. 27th ICRC, Hamburg (2001)/
http://ik1au1.fzk.de/publication/ICRC2001_talk_kampert.pdf /
astro-ph/0204205 (2002).
- [15] D. Heck, J. Knapp, J.N. Capdevielle, G. Schatz, T. Thouw, Forschungszentrum Karlsruhe Report, FZKA 6019 (1998) 90p.
- [16] N.N. Kalmykov, S.S. Ostapchenko, Yad. Fiz. **56** (1993) 105 / Phys.At.Nucl. **56** (3) (1993) 346.
- [17] G. Schatz, "Determining Energy Spectra for Separate Mass Groups from EAS: the Quest Is Still on." Preprint submitted to Elsevier Science (2002).
- [18] S.V. Ter-Antonyan and P.L. Biermann, Preprint, astro-ph/0106076 (2001).
- [19] S.V. Ter-Antonyan, L.S. Haroyan, Preprint, hep-ex/0003006 (2000).
- [20] S.V. Ter-Antonyan and P.L. Biermann, Proc. 27th ICRC Hamburg, HE225 (2001) 1051.